

Hadron Resonances in the Medium via Electromagnetic Probes

Hendrik van Hees

Goethe University Frankfurt

July 07, 2015



FIAS Frankfurt Institute
for Advanced Studies



- 1 Electromagnetic probes and hadron resonances
- 2 Coarse-grained transport approach (UrQMD with Stephan Endres)
- 3 Dielectrons at SIS energies (HADES)
- 4 Dimuons at SPS (NA60)
- 5 Conclusions

Em. current correlator in-medium approaches

Em. current correlation function and dileptons

- dilepton thermal emission rates given by same electromagnetic-current-correlation function ($J_\mu = \sum_f Q_f \bar{\psi}_f \gamma_\mu \psi_f$)

[McLerran+Toimela (1985), Weldon (1990), Gale+Kapusta (1991),...]

$$\Pi_{\mu\nu}^<(q) = \int d^4x \exp(iq \cdot x) \langle J_\mu(0) J_\nu(x) \rangle_T = -2 f_B(q \cdot u) \text{Im} \Pi_{\mu\nu}^{(\text{ret})}(q)$$

$$\frac{dN_{e^+e^-}}{d^4x d^4q} = -g^{\mu\nu} \frac{\alpha^2}{3q^2 \pi^3} \text{Im} \Pi_{\mu\nu}^{(\text{ret})}(q) \Big|_{q^2=M_{e^+e^-}^2} f_B(q \cdot u)$$



- u : four-velocity of the fluid cell; $p \cdot u = p_0^{\text{hb}}$ energy in “heat-bath frame”
- to lowest order in α : $e^2 \Pi_{\mu\nu} \simeq \Sigma_{\mu\nu}^{(\gamma)}$
- vector-meson dominance model:

$$\Sigma_{\mu\nu}^{(\gamma)} = \text{---} \overset{G_\rho}{\text{---}} \text{---}$$

- need in-medium spectral functions of vector bosons
- additional sources: QGP, “multi-pion” processes


Hadronic many-body theory

- HMBT for vector mesons [Ko et al, Chanfray et al, Herrmann et al, Rapp et al, ...]
- $\pi\pi$ interactions and **baryonic excitations**



- +corresponding vertex corrections \Leftrightarrow gauge invariance
- **Baryon (resonances)** important, even at RHIC with low **net** baryon density
 $n_B - n_{\bar{B}}$
- reason: $n_B + n_{\bar{B}}$ relevant (CP inv. of strong interactions)

In-medium spectral functions and baryon effects




[R. Rapp, J. Wambach Adv. Nucl. Phys **25**, 1 (2000), R. Rapp, J. Wambach, HvH, Landolt-Börnstein I/23, 4A.1 (2009), arXiv: 0901.3289 [hep-ph]]

- **baryon effects** important

- large contribution to broadening of the peak
- responsible for most of the strength at small M

In-medium spectral functions and baryon effects




[R. Rapp, J. Wambach Adv. Nucl. Phys **25**, 1 (2000), R. Rapp, J. Wambach, HvH, Landolt-Börnstein I/23, 4A.1 (2009), arXiv: 0901.3289 [hep-ph]]

- **baryon effects** important

- large contribution to broadening of the peak
- responsible for most of the strength at small M

In-medium spectral functions and baryon effects



[R. Rapp, J. Wambach Adv. Nucl. Phys **25**, 1 (2000), R. Rapp, J. Wambach, HvH, Landolt-Börnstein I/23, 4AÅ§1 (2009), arXiv: 0901.3289 [hep-ph]]

- **baryon effects** important

- large contribution to broadening of the peak
- responsible for most of the strength at small M

Coarse-grained UrQMD

(with S. Endres, J. Weil, M. Bleicher)

Coarse-grained UrQMD (CGUrQMD)

- problem with **medium modifications** of spectral functions/interactions
- only available in equilibrium many-body QFT models
- use “in-medium cross sections” naively: **double counting!?**
- way out: map transport to **local-equilibrium fluid**
- use **ensemble of UrQMD** runs with an **equation of state**
- fit **temperature, chemical potentials, flow-velocity field**
from anisotropic energy-momentum tensor [W. Florkowski et al NPA **904-905**, 803c (2013)]


$$T^{\mu\nu} = (\epsilon + P_{\perp})u^{\mu}u^{\nu} - P_{\perp}g^{\mu\nu} - (P_{\parallel} - P_{\perp})V^{\mu}V^{\nu}$$

- thermal rates from **partonic/hadronic QFT become applicable**
- **extrapolated lattice QGP** and **Rapp-Wambach hadronic many-body theory**

Dielectrons (SIS/HADES)


CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- coarse-graining method works at low energies!
- UrQMD-medium evolution + RW-QFT rates [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from Ar + KCl(1.76 AGeV) $\rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]
- $M_{ee} < 0.13$ GeV




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from Ar + KCl(1.76 AGeV) $\rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]
- $0.13 \text{ GeV} < M_{ee} < 0.3 \text{ GeV}$




- dielectron spectra from Ar + KCl(1.76 AGeV) $\rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]
- $0.3 \text{ GeV} < M_{ee} < 0.45 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from Ar + KCl(1.76 AGeV) $\rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]
- $0.45 \text{ GeV} M_{ee} < 0.65 \text{ GeV}$




CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)


- dielectron spectra from Ar + KCl(1.76 AGeV) $\rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]
- $M_{ee} > 0.65$ GeV



CGUrQMD: Ar+KCl (1.76 AGeV) (SIS/HADES)

- dielectron spectra from Ar + KCl(1.76 AGeV) $\rightarrow e^+e^-$ (SIS/HADES)
- m_t spectra [S. Endres, HvH, J. Weil, M. Bleicher, arXiv: 1505.06131 [nucl-th]]
- rapidity spectrum ($M_{ee} < 0.13$ GeV)





- caveat: pp/np acceptance filter with single-e cut, $p_t < 100$ MeV
- correct filter urgently needed!
- excellent agreement with preliminary HADES data
 (data points not shown here on request of the HADES collaboration)


Dimuons (SPS/NA60)

CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- note the importance of **baryon effects!**




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+ \mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- higher IMR: provides **averaged true temperature**
(no blueshifts in the **invariant-mass spectra!**)




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $p_T < 0.2$ GeV




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $0.2 \text{ GeV} < p_T < 0.4 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $0.4 \text{ GeV} < p_T < 0.6 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $0.6 \text{ GeV} < p_T < 0.8 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $0.8 \text{ GeV} < p_T < 1.0 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $1.0 \text{ GeV} < p_T < 1.2 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $1.2 \text{ GeV} < p_T < 1.4 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $1.4 \text{ GeV} < p_T < 1.6 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $1.6 \text{ GeV} < p_T < 1.8 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $1.8 \text{ GeV} < p_T < 2.0 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $2.0 \text{ GeV} < p_T < 2.2 \text{ GeV}$




CGUrQMD: In+In (158 AGeV) (SPS/NA60)

- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+\mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]


- min-bias data ($dN_{\text{ch}}/dy = 120$)
- $2.2 \text{ GeV} < p_T < 2.4 \text{ GeV}$



- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+ \mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]


- min-bias data ($dN_{\text{ch}}/dy = 120$)



- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+ \mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]


- min-bias data ($dN_{\text{ch}}/dy = 120$)



- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+ \mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]


- min-bias data ($dN_{\text{ch}}/dy = 120$)



- dimuon spectra from In + In(158 AGeV) $\rightarrow \mu^+ \mu^-$ (NA60)

[S. Endres, HvH, J. Weil, M. Bleicher, PRC **91**, 054911 (2015)]

- min-bias data ($dN_{\text{ch}}/dy = 120$)



- em. probes, $\ell^+\ell^-$ and γ : **negligible final-state interactions**
- probe **in-medium electromagnetic current-current correlator** over **entire history of fireball evolution**
- provide insight into fundamental properties of **QCD matter**
- needs models for electromagnetic radiation from **QGP and hadron gas**
- medium effects on **vector mesons in hot and dense matter**
- hint at **chiral-symmetry restoration**
⇒ melting resonances rather than dropping mass
- prevalence of **baryon resonance interactions** in vector-meson SFs!