

Clusters and Hypernuclei formation study within PHQMD Model

V. Kireyeu¹, A. Le Fèvre², E. Bratkovskaya², J. Aichelin³, Y. Lefeils²


XXIVth International Baldin Seminar on High Energy Physics Problems
21.09.2018

1 - JINR, Dubna, Russia

2 - GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

3 - SUBATECH, UMR 6457, Ecole des Mines de Nantes - IN2P3/CNRS - Université de Nantes, France

Introduction




At 3 A.GeV even in central collisions almost 20% of the baryons are bound in the clusters

Without dynamical fragments formation we cannot properly describe observables like v_1 , v_2 , p_T spectra,

(it's a lot!)

20%



Many present transport models fail to describe fragments at NICA/FAIR (and higher) energies. We made a new one.

PHSD

E.L. Bratkovskaya, W. Cassing, Nucl.Phys. A856 (2011) 162-182.



Initial A+A collisions - HSD: string formation and decay to pre-hadrons

Fragmentation of pre-hadrons into quarks: using the quark spectral functions from the Dynamical QuasiParticle Model (DQPM) approximation to QCD

DQPM: Peshier, Cassing, PRL 94 (2005) 172301; Cassing, NPA 791 (2007) 365; NPA 793 (2007)

Partonic phase: quarks and gluons (= „dynamical quasiparticles“) with off-shell spectral functions (width, mass) defined by DQPM

elastic and inelastic parton-parton interactions:
using the effective cross sections from the DQPM

- ✓ $q + q\bar{q}$ (flavor neutral) \Leftrightarrow gluon (colored)
- ✓ gluon + gluon \Leftrightarrow gluon (possible due to large spectral width)
- ✓ $q + q\bar{q}$ (color neutral) \Leftrightarrow hadron resonances

Hadronization: based on DQPM - massive, off-shell quarks and gluons with broad spectral functions hadronize to off-shell mesons and baryons:

gluons \rightarrow $q + q\bar{q}$; $q + q\bar{q} \rightarrow$ meson (or string);
 $q + q + q \rightarrow$ baryon(or string)(strings act as ‚doorway states‘ for hadrons)

Hadronic phase: hadron-string interactions - off-shell HSD


FRIGA

A. Le Fèvre et al., J. Phys.: Conf. Ser. 668 (2016) 012021.


1) Pre-select good «candidates» for fragments according to proximity criteria: real space coalescence = **Minimum Spanning Tree (MST)** procedure.

2) Take randomly 1 nucleon out of one fragment


3) Add it randomly to another fragment



$$E = E_{kin}^1 + E_{kin}^2 + V^1 + V^2$$



$$E' = E_{kin}^{1'} + E_{kin}^{2'} + V^{1'} + V^{2'}$$



If $E' < E$ take the new configuration

If $E' > E$ take the old with a probability depending on $E' - E$

Repeat this procedure very many times...


It leads automatically to the most bound configuration.

PHQMD

Parton-Hadron Quantum Molecular Dynamics
= PHSD + QMD* + FRIGA

* J. Aichelin and H. Stöcker, Phys. Lett. 176 B (1988) 14


Model predictions



Single particle spectra still the same as in PHSD

Produced particles are well reproduced at NICA/FAIR/SPS energies



Model predictions



Single particle spectra still the same as in PHSD

Produced particles are well reproduced at NICA/FAIR/SPS energies

Model predictions





Single particle spectra still the same as in PHSD

Produced particles are well reproduced at NICA/FAIR/SPS energies

Model predictions


(preliminary results at NICA energies)



Central collisions: light clusters;
Semi-peripheral collisions: existence of heavy clusters – remnants from spectators

Clusterization time

Maybe here..?







Oops...

TOO LATE





Clusterization time



M_{imf} vs Z_{bound} @ 1.23 GeV

Courtesy of the ALADIN Collaboration for the new S254 data

Au+Au, $E_{\text{lab}} = 1.23$ A.GeV, $t_{\text{pass}} = 16.86$ fm/c





$\langle M_{\text{IMF}} \rangle$ – average number of medium mass fragments ($2 < Z < 30$)

Z_{bound} – number of charges bounded in clusters ($Z > 1$)

M_{imf} vs Z_{bound} @ 2 GeV

Courtesy of the ALADIN Collaboration for the new S254 data

Au+Au, $E_{\text{lab}} = 2.00$ A.GeV, $t_{\text{pass}} = 13.22$ fm/c





$\langle M_{\text{IMF}} \rangle$ – average number of medium mass fragments ($2 < Z < 30$)

Z_{bound} – number of charges bounded in clusters ($Z > 1$)

M_{imf} vs Z_{bound} @ 4 A.GeV

Courtesy of the ALADIN Collaboration for the new S254 data

Au+Au, $E_{\text{lab}} = 4.00$ A.GeV, $t_{\text{pass}} = 9.35$ fm/c





$\langle M_{\text{IMF}} \rangle$ – average number of medium mass fragments ($2 < Z < 30$)

Z_{bound} – number of charges bounded in clusters ($Z > 1$)

M_{imf} vs Z_{bound} @ 6 A.GeV

Courtesy of the ALADIN Collaboration for the new S254 data

Au+Au, $E_{\text{lab}} = 6.00$ A.GeV, $t_{\text{pass}} = 7.63$ fm/c





$\langle M_{\text{IMF}} \rangle$ – average number of medium mass fragments ($2 < Z < 30$)

Z_{bound} – number of charges bounded in clusters ($Z > 1$)

M_{imf} vs Z_{bound} @ 8 A.GeV

Courtesy of the ALADIN Collaboration for the new S254 data

Au+Au, $E_{\text{lab}} = 8.00$ A.GeV, $t_{\text{pass}} = 6.61$ fm/c



$\langle M_{\text{IMF}} \rangle$ – average number of medium mass fragments ($2 < Z < 30$)

Z_{bound} – number of charges bounded in clusters ($Z > 1$)

M_{imf} vs Z_{bound} @ 11 A.GeV



Courtesy of the ALADIN Collaboration for the new S254 data

$\sqrt{s_{NN}} \approx 5$ GeV

NICA!

Au+Au, $E_{lab} = 11.00$ A.GeV, $t_{pass} = 5.64$ fm/c

Au+Au, $E_{lab} = 11.00$ GeV, $t_{pass} = 5.64$ fm/c




$\langle M_{IMF} \rangle$ – average number of medium mass fragments ($2 < Z < 30$)

Z_{bound} – number of charges bounded in clusters ($Z > 1$)

Why not to use just coalescence?

Courtesy of the ALADIN Collaboration for the new S254 data

Au+Au, $E_{\text{lab}} = 11.00$ GeV, $t_{\text{pass}} = 5.64$ fm/c




It fails to describe spectators

Not very healthy for the Flow Analysis


PHQMD+FRIGA
may be also used for engineering stuff

We can estimate damage caused to detector


Z vs Θ @ 1.23 A.GeV



Z vs Θ @ 2 A.GeV



Z vs Θ @ 4 A.GeV



Z vs Θ @ 4 A.GeV

Au+Au, $E_{\text{lab}} = 4.00$ A.GeV




Summary

- PHQMD can produce clusters and hypernuclei;
- Model reproduce experimental data;
- Model`s predictions can be used for analysis, feasibility and engineering studies;
- Model is actively developing.

Backup


Old

Au+Au, $\sqrt{s} = 5$ GeV, $b = 0.3$ fm



New


AuAu, $\sqrt{s} = 5$ GeV, $b = 0.3$ fm



Backup


Fragments $Z \geq 2$

Au+Au, $E_{\text{lab}} = 11.00$ GeV, $t_{\text{pass}} = 5.64$ fm/c



Hypernuclei

Au+Au, $E_{\text{lab}} = 11.00$ GeV, $t_{\text{pass}} = 5.64$ fm/c



Backup

